Space Weather Technical Forecast

Issued on Wednesday, 27 December 2023 at 12:40 Local

This technical forecast provides a four day assessment of space weather events. The probabilities stated below are for reaching or exceeding the given levels. For more information about space weather impacts please see the Met Office Space Weather Scales https://www.metoffice.gov.uk/weather/learn-about/space-weather/uk-scales

Space Weather Forecast Headline: Risk of G1/Minor Storm or perhaps peak G2/Moderate Storm from a series of CMEs on days 1 and 2 (27th and 28th Dec). Slight Chance of further R1 Radio Blackouts.

Analysis of Space Weather Activity over past 24 hours

Solar Activity: Solar activity, as measured by GOES16, has been Low over the past 24 hours, with only minor C-class flares observed.
There are currently five numbered sunspot regions and one unnumbered region on the visible disc, the unnumbered region is showing signs of growth on the southeast limb, but remains small and simple at present. AR3530 in the northeast has shown some minor growth since previous analysis. AR3521 is due to rotate off around western limb soon, AR3526 has faded to plage. All remaining regions appear largely unchanged or in slow decay. No new Earth-directed Coronal Mass Ejections (CMEs) were seen in imagery, although numerous potential sources remain in the form of many filaments on the disc.
Solar Wind / Geomagnetic Activity: Solar winds, as measured by ACE and DSCOVR at L1, were mostly Slightly Elevated throughout the period, briefly reaching Elevated levels, with a peak solar wind speed of 538km/s at 26/1913 UTC. The solar wind density was average and just below 10ppcc throughout the period. The IMF was weak at 6nT or lower, while the all-important north-south component was variable and weak. The phi angle was mostly positive (away from the Sun). The net result of the above solar wind measures was for provisionally Quiet geomagnetic activity throughout the period.
Energetic Particles / Solar Radiation: GOES16 high energy (greater than 10MeV) proton flux was very slightly raised above Normal Background levels in the 24 hours, although well below S1 (10pfu).
GOES16 high energy (greater than 2MeV) electron flux was mostly at Normal Background levels. The 24-hour fluence was below Active throughout and declining, with the 27/0001UTC observation being 4.86e6 integrated pfu.

Issued by Met Office Space Weather Advisor, Tel: +44 (0) 330 135 4254 Email: moswoc@metoffice.gov.uk

© Crown Copyright 2023. All Rights Reserved.

Met Office Space Weather Technical Forecast

Four-Day Space Weather Forecast Summary

Solar Activity: Low to Moderate activity is forecast, with a slight chance of isolated M-class X-ray flares.

Solar Wind / Geomagnetic Activity: There has been a glut of CMEs over the Christmas period, with the distilled result being of two possible glancing blows and one more assured hit. These events are likely to interact and as such there is lower than average confidence in the geomagnetic forecast, even though no coronal hole fast winds feature in the forecast.

A pair of CME glancing blows are expected on Days 1 and 2 (27 and 28 December), with a third more direct hit also possible on 28 December. The possibility that this final CME arrives concurrently with the second glancing blow should give rise to a peak risk of G1/Minor Storm, and perhaps even G2 in the UTC day of 28 December, with a residual tail of effects perhaps continuing into 29 December. Geomagnetic activity should return to mostly Quiet conditions by Day 4 (30 December).

Energetic Particles / Solar Radiation: GOES16 high energy proton flux is near Background and likely to persist at this level, but with a Slight Chance of increasing given any further notable flares from AR3529 or other regions on the disc.

GOES16 high energy electron flux is expected to persist at mainly Normal Background levels, with the associated fluence remaining below the Active level throughout.

Issued by Met Office Space Weather Advisor, Tel: +44 (0) 330 135 4254 Email: moswoc@metoffice.gov.uk

© Crown Copyright 2023. All Rights Reserved.

Figure 1. Solar Analysis Valid 27/0800UTC.



.

Key: Filament ___ , Prominence ___ , Magnetic Field Line - - - , Polarity +/-, Coronal Holes: Grey shaded area CHxx +/-, Sunspot groups 25xx - Mt Wilson α-β-βγ-βγδ and Zurich-McIntosh Axx etc.

Issued by Met Office Space Weather Advisor, Tel: +44 (0) 330 135 4254 Email: moswoc@metoffice.gov.uk

© Crown Copyright 2023. All Rights Reserved.

Met Office Space Weather Technical Forecast

Geomagnetic Storms:

No coronal hole fast winds feature in the forecast.
A great number of CMEs were observed over the Christmas period, however three have stood out as being of potentially more relevance. Overall, the forecast carries high uncertainty given the marginal effects from two of these three CMEs, but also due to their probable inter-connectivity, with each possibly priming the interplanetary medium for the next. The most noteworthy events are summarised as:

Firstly, a large filament eruption from the northwest disc around 24/1200UTC. This may give a glancing blow from around the middle of the UTC day of 27 December, but is aimed mainly above and ahead of the Sun-Earth line.
Secondly, a coronal dimming from the southwest disc, around 24/1430UTC. This may deliver a second glancing blow around UTC dawn on 28 December. This CME is angled mainly below and ahead of the Sun-Earth line.
Thirdly, and most directly, a coronal dimming that occurred just northeast of centre-disc around 25/0700UTC. This presented on coronagraph imagery as a halo with a slight eastward bias. This will likely affect Earth, however the main uncertainty here is the arrival time - given the poor triangulation offered by STEREO and Lasco viewpoints at this stage of the cycle, but also because the earlier glancing CMEs may impede or sweep clear the interplanetary medium ahead of this following halo.
MOSWOC Enlil has come close at times to modelling these three CMEs as a very large united density front, but more frequently has kept them as closely-spaced separate events. The model itself appears to have a reasonable treatment of current winds and forecast ambient conditions of fast winds, which increases confidence in CME transit times to 1AU, assuming the analyses were reasonably accurate.
Geomagnetic activity will be most likely Quiet at first though will increase should Earth receive any contact from the CMEs. 27/1200UTC (Day 1) and 28/0600UTC (Day 2) are the most likely times for the glancing blows to pass 1AU, giving a Chance of G1/Minor Storm. It is possible that Earth could 'thread the needle' and see both of these events pass to no effect, however the third CME appears unavoidable, with a most likely arrival time of 28/1400UTC (Day 2), but with very large error bars (perhaps +/-12 hours). Although weak in itself, this third CME may therefore arrive in close proximity to one of the glancing blows, which will in any case lead to the peak risk of geomagnetic activity in the period, and perhaps even a Moderate Storm G2, although G1 is much more Likely. A tail of risk then extends into Day 3 (29 December), becoming conditions return to mostly Quiet on Day 4 (30 December).

Geo-Magnetic Storm Probability (Exceedance)	Level	Past 24 Hours (Yes/No)	Day 1 (00-24 UTC) (%)	Day 2 (00-24 UTC) (%)	Day 3 (00-24 UTC) (%)	Day 4 (00-24 UTC) (%)
Minor or Moderate	G1 to G2	No	40	60	20	1
Strong	G3	No	1	5	1	1
Severe	G4	No	1	1	1	1
Extreme	G5	No	1	1	1	1

Geomagnetic Activity - Earthbound Coronal Mass Ejections

Issued by Met Office Space Weather Advisor, Tel: +44 (0) 330 135 4254 Email: moswoc@metoffice.gov.uk

© Crown Copyright 2023. All Rights Reserved.

Date/time 21.5R (UTC)	Halo: Full or Partial	Source	Source Location	Estimated Speed	Estimated Arrival Time	Comments
24/1732	Partial	Filament	NW	850km/s	27/1200UTC +/-12 hours	Low confidence, glancing blow
24/2224	Partial	Filament	SW	572km/s	28/0600UTC +/-12 hours	Low confidence, glancing blow
25/0700	Full halo	Filament	NE centre	538km/s	28/1400UTC +/-12 hours	High confidence hit, low confidence timing and interaction with other CMEs

Figure 2: MOSWOC Enlil with potentially up to the Earth-Directed CMEs



Issued by Met Office Space Weather Advisor, Tel: +44 (0) 330 135 4254 Email: moswoc@metoffice.gov.uk

© Crown Copyright 2023. All Rights Reserved.

Radio Blackouts - X-Ray Flares:
Low to Moderate activity is forecast, with a daily Slight Chance of isolated M-class flares.
The X-ray trace has settled near the B/C flux boundary, with occasional low C-class activity. The sunspots on the facing side show relatively little complexity at present and are either stable and mature, or else slowly fading. AR3530 developed some intermediate spots and the NEW region in the southeast will need monitoring.
An M-class flare was observed over the southeast limb on 24 December, and it remains to be seen if this is the precursor to a more active region rotating onto the disc. Along with a distinct area of magnetic looping visible on AIA171 in the northeast, this may suggest the potential for further arriving regions, although these will be assessed when they're present for analysis.

X Ray Flares	Level	Past 24 Hours (Yes/No)	Day 1 (00-24 UTC)	Day 2 (00-24 UTC)	Day 3 (00-24 UTC)	Day 4 (00-24 UTC)
Probability			(%)	(%)	(%)	(%)
Active	R1-R2 M Class	No	20	20	20	20
Very Active	R3 to R5 X Class	No	1	1	1	1

Figure 3: GOES18 X-Ray flares



Issued by Met Office Space Weather Advisor, Tel: +44 (0) 330 135 4254 Email: moswoc@metoffice.gov.uk

© Crown Copyright 2023. All Rights Reserved.

Solar Radiation Storms - (High Energy Protons):

GOES16 high energy (greater than 10MeV) proton flux was very slightly raised above resting Background levels in the 24 hours, although below S1 (10pfu) and in fact even below 1pfu for the duration. The cause of this enhancement is not known, however given the very gradual rise and soft spectrum, it may be a far side or even eastern limb event, perhaps even the long-period M1.1 from behind the eastern horizon on 24 December.
There is a daily Slight Chance of proton enhancement if any further M-class flare activity were to occur.

Radiation Storms Probability (Exceedance)	Level (cm^{-2} sr^{-1} s^{-1})	Past 24 Hours (Yes/No)	Day 1 (00-24 UTC) (%)	Day 2 (00-24 UTC) (%)	Day 3 (00-24 UTC) (%)	Day 4 (00-24 UTC) (%)
Active	**≥ S1**	No	5	5	5	5
Very Active	**≥ S3 ***	No	1	1	1	1
* S3 ≥ 10 MeV ≥ 1000 pfu and / or ≥ 50 MeV ≥ 10 pfu. (pfu = $cm^{-2}sr^{-1}s^{-1}$)						

High Energy Electrons Event (≥ 2MeV):

High energy electron flux (greater than 2MeV) observed by GOES16 is mostly at Normal Background, with no further coronal hole influences featured in the forecast, although brief periods of Moderate flux are likely. There are then the CME glancing impacts possible from 27 to 29 December, and these are likely to further suppress observed flux at GEO until after this four-day period.
The associated 24-hour fluence is expected to persist below Active (1e8 pfu) as a result of the above. Regarding MOSWOC REFM, the fact that the current possible fast wind enhancement is not forecast nor a persistent feature, as well as the presence of three possible CMEs later in the period, will mean that although the initial trend is accepted, the rational is probably incorrect. MOSWOC REFM should not be followed in the second half of the period, although the forecast low fluence count is probably correct.

GEO Electron Environment Probability (Exceedance)	Level (cm^{-2} sr^{-1} day^{-1})	Past 24 Hours (Yes/No)	Day 1 (00-24 UTC) (%)	Day 2 (00-24 UTC) (%)	Day 3 (00-24 UTC) (%)	Day 4 (00-24 UTC) (%)
Active	**≥ 2 MeV ≥ $1x10^8$**	No	1	1	1	5
Very Active	**≥ 2 MeV ≥ $1x10^9$**	No	1	1	1	1

Issued by Met Office Space Weather Advisor, Tel: +44 (0) 330 135 4254 Email: moswoc@metoffice.gov.uk

© Crown Copyright 2023. All Rights Reserved.

Figure 4: MOSWOC REF Model



Issued by Met Office Space Weather Advisor, Tel: +44 (0) 330 135 4254 Email: moswoc@metoffice.gov.uk

© Crown Copyright 2023. All Rights Reserved.